

PANTONE 374c
PANTONE 374c 30%
M100%
C100% M100%
K100%

複方15x8.3cm

Ingredient(s): Each 6.0 g contains the extracts equivalent to raw herb

成分：每6.0公克中含提取物相当于原生药

柴胡 Bupleuri Radix..... 6.0g
制半夏 Pinelliae Rhizoma Praeparata..... 3.8g
黄芩 Scutellariae Radix..... 2.3g
白芍 Paeoniae Alba Radix..... 2.3g
大枣 Jujubae Fructus..... 2.3g
枳实 Aurantii Immaturus Fructus..... 2.0g
大黄 Rhei Radix et Rhizoma..... 1.5g
生姜 Zingiberis Rhizoma Recens..... 3.8g
以上生药制成浸膏..... 4.0g

(生药与浸膏比例24:4=6:1)

许可证字号：台湾卫署药制字第028638号

Batch No.:

Expiry Date:



“勝昌”

203090

大柴胡汤

浓缩散

Da Chai Hu Tang

Extract Pulveres “Sheng Chang”

【中药浓缩制剂】

重量Net Wt.: 200 g

Manufacturer :
SHENG CHANG PHARMACEUTICAL CO., LTD.
ZHONGLI FACTORY (G.M.P.) Taoyuan, Taiwan
Importer :
LONG CHANG HERBAL (S) PTE LTD
196 Pandan Loop #05-16
Pantech Business Hub Singapore 128384

Indication(s):

Cold damage entering yang brightness, incompletely release the lesser yang disease, alternating chills and fever, stuffiness and fullness below the heart, persistent vomiting, vexation and constipation.

适应症:

伤寒入阳明、少阳未尽。寒热往来、心下鞕满、呕吐不止、心煩便秘。

Dosage:

For Adults: 2.0 g each time, 3 times per day before meal with luke-warm water.

Children: The dose is gradually decreased by age.

Use only as directed by physician.

用法用量:

成人每次服用2.0公克，每日3次，在饭前一小时以温开水送服。儿童依年龄递减之。本药须由中医师处方使用。

Contraindication(s): Contraindicated in syndrome of yin deficiency accompanied with dry cough, body fluid depletion and thirst, in hemorrhagic disorders and in dry-phlegm syndrome.

Use with caution in pregnancy, lactation, menstruation, children and undiagnosed abdominal symptoms such as pain, nausea or vomiting. Use with caution in individuals with cold deficiencies in spleen and stomach or with qi and blood deficiencies. Avoid prolonged use.

禁忌: 阴虚燥咳、津伤口渴、血证及痰燥者禁服。孕妇、哺乳期妇女、月经期妇女、儿童以及未确诊的腹痛、恶心、呕吐患者慎用。脾胃虚寒、血虚气弱者慎用。避免长期服用。

Side effects: May cause nausea, vomiting and cramp-like discomfort of the gastrointestinal tract; long-term use may lead to electrolyte disturbances and aggravation of constipation with dependency.

副作用: 可能引起恶心、呕吐、腹痛等症狀。长期服用可能会引起电解质紊乱及产生依赖性便秘。

Allowed for sale as a Chinese Proprietary Medicine based on information submitted to the Authority. Consumer discretion is advised.
根据向当局提交的资料允许作为中成药销售。
谨慎选用。