複方15x8.3cm

Ingredient(s): Each 7.5 g contains the extracts equivalent to raw herb

成	分:	每7.5公克中含提取物相当于原	生药
麻	1	Ephedrae Herba	. 5.0g
炙	甘草	🕏 Glycyrrhizae Radix Praeparata	10.0g
薏	苡仁	Coicis Semen	. 5.0g
杏	4	Armeniacae Amarum Semen	. 4.0g
以	上生	上药制成浸膏	. 3.4g
(生	药-	与浸膏比例24:3.4=7.1:1)	

许可证字号:台湾卫署药制字第028428号

Batch No.: **Expiry Date:**





"勝昌"

211290

麻杏薏甘汤

Ma Xing Yi Gan Tang

Extract Powder "Sheng Chang"

【中药浓缩制剂】 重量Net Wt.: 200 g

Manufacturer :

SHENG CHANG PHARMACEUTICAL CO., LTD. ZHONGLI FACTORY (G.M.P.) Taovuan. Taiwan Importer :

LONG CHANG HERBAL (S) PTE LTD 196 Pandan Loop #05-16 Pantech Business Hub Singapore 128384 Indication(s):

Wind-dampness, generalized pain and fever, which worse in the late afternoon.

风湿身痛发热,日晡甚者。

For Adults: 2.5 g each time, 3~4 times per day before meal with

Children: The dose is gradually decreased by age. Use only as directed by physician.

成人每次服用2.5公克,每日3~4次,在饭前一小时以 温开水送服。儿童依年龄递减之 本药须由中医师处方使用。

Contraindication(s): Contraindicated in individuals with loose stools, cough due to yin-deficiency, weak constitution, spontaneous perspiration, night sweat or dyspnea of asthenia type. Use with caution in babies. Avoid overdose. Recommend advice from physicians prior to consumption in babies. Use with caution in pregnancy, lactation and individuals with heart diseases, hypertension, glaucoma, diabetes and thyroid diseases. Avoid prolonged use.

禁忌:阴虚咳嗽、大便溏泻、体虚自汗、盗汗及虚喘者 禁服。嬰儿慎服。不宜过量服用。孕妇、哺乳期妇女、 心脏病、高血压、青光眼、糖尿病以及甲状腺疾病患者 慎用。避免长期服用。

Side effects: May cause dizziness, headaches, poor appetite, insomnia, irregular or fast heartbeat and rise in blood pressure.

副作用:可能会引起头晕、头痛、食欲不振、失眠、心 悸或心律加快以及血压升高。

Allowed for sale as a Chinese Proprietary Medicine based on information submitted to the Authority. Consumer discretion is advised. 根据向当局提呈的资料允许作为中成药销售 谨慎选用